## THE STATIONARITY OF AN ESTIMATED AUTOREGRESSIVE PROCESS

BY T. W. ANDERSON

TECHNICAL REPORT NO. 7
NOVEMBER 15, 1971

PREPARED UNDER CONTRACT
N00014-67-A-0112-0030 (NR-042-034)
FOR THE OFFICE OF NAVAL RESEARCH
THEODORE W. ANDERSON, PROJECT DIRECTOR

DEPARTMENT OF STATISTICS
STANFORD UNIVERSITY
STANFORD, CALIFORNIA



## THE STATIONARITY OF AN ESTIMATED AUTOREGRESSIVE PROCESS

by

T. W. Anderson Stanford University

TECHNICAL REPORT NO. 7

NOVEMBER 15, 1971

PREPARED UNDER THE AUSPICES OF
OFFICE OF NAVAL RESEARCH CONTRACT #N00014-67-A-0112-0030
T. W. ANDERSON, PROJECT DIRECTOR

DEPARTMENT OF STATISTICS
STANFORD UNIVERSITY
STANFORD, CALIFORNIA

A stationary stochastic process  $\{y_t\}$  with mean  $\{y_t\} = 0$  satisfies a stochastic difference equation if there exist constants  $\beta_0=1,\ \beta_1,\ \ldots,\ \beta_r$  such that  $\{u_t\}$  defined by

(1) 
$$\sum_{r=0}^{k} \beta_r y_{t-r} = u_t, \qquad t = \dots, -1, 0, 1, \dots,$$

consists of independently identically distributed random variables. The process  $\{y_t^{}\}$  is stationary and  $y_t^{}$  is independent of  $u_{t+1}^{}$ ,  $u_{t+2}^{}$ , ... if and only if the roots of the associated polynomial equation

(2) 
$$\sum_{r=0}^{p} \beta_r x^{p-r} = 0$$

are less than 1 in absolute value. The process is autoregressive of order p. We assume  $\xi u_t = 0$  and  $\xi u_t^2 = \sigma^2$  with  $0 < \sigma^2 < \infty$ .

Let  $y_1, \ldots, y_T$  be T successive observations on the process. To estimate the coefficients  $\beta_1, \ldots, \beta_p$  one can solve the linear equations

(3) 
$$\sum_{j=1}^{p} c_{j-j} b_{j} = -c_{j}, \qquad j=1, ..., p,$$

where

(4) 
$$c_i = c_{-i} = \frac{1}{T} \sum_{t=1}^{T-i} y_{t+i} y_t, \quad i=0, 1, ..., p.$$

See, for example, Section 5.4 of T. W. Anderson (1971). We assume that there are at least p different nonzero values of t observed. The purpose of this note is to show that the solution of (3) yields coefficients corresponding to a stationary process; that is, the roots of

(5) 
$$\sum_{r=0}^{p} b_r x^{p-r} = 0$$

are less than 1 in absolute value. Pagano (1971) has shown this result by a different method.

Let  $y_{-p+1} = y_{-p+2} = \dots = y_0 = 0$  and  $y_{T+1} = y_{T+2} = \dots = y_{T+p} = 0$ . Define the vectors

(6) 
$$\widetilde{y}_{t} = \begin{pmatrix} y_{t} \\ y_{t-1} \\ \vdots \\ y_{t-p+1} \end{pmatrix}, \qquad t=0, 1, \dots, T+p.$$

The equations (3) can be written

The equation (7) is the first row of

(8) 
$$\overset{\mathbf{T}+\mathbf{p}}{\underset{t=1}{\Sigma}} \quad \overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}}{\overset{\mathbf{\widetilde{y}}_{t}}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}}{\overset{\mathbf{\widetilde{y}}_{t}}{\overset{\mathbf{\widetilde{y}}_{t}}}{\overset{\mathbf{\widetilde{y}}_{t}}}{\overset{\mathbf{\widetilde{y}}}{\overset{\mathbf{\widetilde{y}}}}{\overset{\mathbf{\widetilde{y}}}{\overset{1}}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}}{\overset{1}}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}{\overset{1}}}{\overset{1}}{\overset{1}}$$

and b' is the first row of B. The other p-1 rows of B constitute the matrix (-I 0).

Theorem 1. The matrix  $\tilde{\mathbb{B}}$  defined by (8) has characteristic roots less than 1 in absolute value.

 $\underline{\underline{Proof}}$ . If  $\underline{\underline{u}}$  is a characteristic vector of  $\underline{\underline{b}}$  corresponding to a characteristic root  $\lambda$ 

Normalize u so that

(10) 
$$1 = \underline{u}' \quad \sum_{t=1}^{T+p} \quad \widetilde{y}_t \quad \widetilde{y}_t' \quad \overline{u} = \sum_{t=1}^{T+p} \quad (\underline{u}' \quad \widetilde{y}_t) \quad \overline{(\underline{u}' \quad \underline{y}_t)}$$
$$= \underline{u}' \quad \sum_{t=1}^{T+p} \quad \widetilde{y}_{t-1} \quad \widetilde{y}_{t-1}' \quad \overline{u} = \sum_{t=1}^{T+p} \quad (\underline{u}' \quad \widetilde{y}_{t-1}) \quad \overline{(\underline{u}' \quad \underline{y}_{t-1})} ,$$

where  $\overline{\underline{u}}$  is the complex conjugate of  $\underline{u}$ . Then multiplication of (9) on the right by  $\overline{\underline{u}}$  gives

(11) 
$$\lambda = \underbrace{\mathbf{u}'}_{t=1} \quad \underbrace{\sum_{t=1}^{T+p}}_{t=1} \quad \underbrace{\widetilde{\mathbf{y}}_{t}}_{t=1} \quad \underbrace{\widetilde{\mathbf{u}}}_{u} = - \quad \underbrace{\sum_{t=1}^{T+p}}_{t=1} \quad (\underbrace{\mathbf{u}'}_{u} \quad \widetilde{\mathbf{y}}_{t}) \quad (\overline{\mathbf{u}'}_{u} \quad \widetilde{\mathbf{y}}_{t-1}).$$

By the Cauchy-Schwarz Inequality  $|\lambda| \le 1$ . We can have equality only if  $\underline{u}' \ \tilde{\underline{y}}_t = \underline{u}' \ \tilde{\underline{y}}_{t-1}$ , t=1, ..., T+p, which is impossible. Q.E.D.

Since the characteristic roots of B are the roots of (5), the desired result has been proved. [See Section 5.4 of T. W. Anderson (1971).]

Theorem 2. The roots of (5), where b<sub>1</sub>, ..., b<sub>p</sub> are the solution to (3), are less than 1 in absolute value.

The result can be extended to the vector-valued autoregressive process  $\{y_t\}$  satisfying

(12) 
$$\sum_{r=0}^{p} \ \sum_{r=r}^{p} \ y_{t-r} = u_{t}, \qquad t = \dots, -1, 0, 1, \dots,$$

where  $y_t$  and  $u_t$  are q-component vectors and  $\beta_0 = I$ ,  $\beta_1$ , ...,  $\beta_p$  are  $q \times q$  matrices,  $\xi u_t = 0$ , and  $\xi u_t u_t' = \Sigma$ , positive definite and finite. The analogue of (2) is

$$\left|\sum_{r=0}^{p} \ \tilde{g}_{r} \lambda^{p-r}\right| = 0.$$

We observe  $y_1, \dots, y_T$ , and define

(14) 
$$c_{i} = c'_{-i} = \frac{1}{T} \sum_{t=1}^{T-i} y_{t+i} y'_{t}, \quad i=0, 1, ..., p.$$

Then the estimates  $_{\sim 1}^{B}$ , ...,  $_{\sim p}^{B}$  are the solution to

(15) 
$$\sum_{j=1}^{p} B_{j} c_{i-j} = -c_{i}, \qquad i=1, \ldots, p.$$

The roots of (13) with  $B_r$  replaced by  $B_r$ , r=1, ..., p, have roots less than 1 in absolute value.

## REFERENCES

- Anderson, T. W. (1971), The Statistical Analysis of Time Series, John Wiley & Sons, Inc.
- Pagano, Marcello (1971), "When is an Autoregressive Scheme Stationary?", Research Report No.53, Department of Statistics, State University of New York at Buffalo.

Security Classification

Socurity classification of title, body of abstract and index	HTROL DATA	R&D é entored when	the overall report to classified)			
1. ORIGINATING ACTIVITY (Corporate cuther)  DEPARTMENT OF STATISTICS  STANFORD UNIVERSITY, CALIF.		Se. REPORT SECURITY CLASSIFICATION				
		24 enem	24 GROUP			
3. REPORT TITLE THE STATIONARITY OF AN ESTIMATED AU	TOREGRESSIVE	PROCESS				
4. DESCRIPTIVE NOTES (Type of report and Includive delea) TECHNICAL REPORT						
5. AUTHOR(5) (Lest name, limit name, initial) ANDERSON, T.W.						
6. REPORT DATE November 15, 1971	70. 70 TAL NO. 0	FPAGES	75. NO. OF REFS			
84. CONTRACT OR GRANT NO.  N00014-67-A-0112-0030  b. project no.  NR-042-034	3. ORIGINATOR'S REPORT NUMBER(S)					
·c.	36. OTHER REFOI	rt No(S) (Any	ether numbers that may be assigned			
Reproduction in whole or in part is UNITED STATES GOVERNMENT	s permitted fo	or any pur	pose of the			
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY OFFICE OF NAVAL RESEARCH ARLINGTON, VIRGINIA					
13. ABSTRACT,						

The estimated coefficients of an autoregressive process define a stationary process if they are computed from the (Toeplitz) matrix of sample moments computed from all available observations, using the same divisor.

Security Classification	LINKA		LINKB		LINKC		
· .	KEY WORDS	ROLE	WT	ROLE	WT	ROLE.	94 L
BANK LISE YERRING SITTE SOFFERENCE MAN	THE SECRET OF THE PROPERTY AND SECRET SECRET AND SECRET				, 1 3 h		
	autoregressive process						
	stationary						
	estimated coefficients						
	associated polynomial equation						
	vector process						
		war a sa					

## INSTRUCTIONS

- ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantse, Department of Dotages activity or other organization (corporate author) issuing the report.
- 2a. REPORT SECURITY CLASSIFICATION: Enter the overall security cleasification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.
- 25. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.
- 3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.
- 4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.
- 5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter test name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.
- 6. REPORT DATE: Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.
- 7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 7b. NUMBER OF REFERENCES: Enter the total number of references cited in the report.
- 8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.
- 8b, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.
- 9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.
- 9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).
- 10. AVAE ABILITY/LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements auch as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and disnemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

- 11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.
- 12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address,
- 13. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shail end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, reles, and weights is optional.